arXiv:quant-ph/0211120v1 19 Nov 2002

Two-photon imaging and quantum holography

Gunnar Björk* and Jonas Söderholm
Department of Microelectronics and Information Technology,
Royal Institute of Technology (KTH), Electrum 229, SE-164 40 Kista, Sweden

Luis L. Sánchez-Soto
Departamento de Óptica, Facultad de Ciencias Físicas, Universidad Complutense, 28040 Madrid, Spain
(Dated: February 1, 2008)

It has been claimed that "the use of entangled photons in an imaging system can exhibit effects that cannot be mimicked by any other two-photon source, whatever strength of the correlations between the two photons" [A. F. Abouraddy, B. E. A. Saleh, A. V. Sergienko, and M. C. Teich, Phys. Rev. Lett. **87**, 123602 (2001)]. While we believe that the cited statement is true, we show that the method proposed in that paper, with "bucket detection" of one of the photons, will give identical results for entangled states as for appropriately prepared classically correlated states.

PACS numbers: 42.50.Dv, 42.30.Lr, 42.30.Va, 03.65.Ta

Recently, there has been a constant focus of interest in quantum limited imaging [1, 2, 3] and lithography [4, 5, 6, 7]. In the former of the applications, the quantum correlation between the amplitude fluctuations of an appropriately entangled multimode state of light is used to beat the standard quantum limit of spatial resolution. The latter application capitalizes on the short de Broglie wavelength of entangled multiphoton states and offers, in principle, a spatial resolution that is independent of the classical wavelength of the electromagnetic field [8].

In a recent paper by Abouraddy *et al.* [9], a method has been proposed where an entangled photon pair is used to image an object. The setup is depicted in Fig. 1. Each of the photons in a photon pair is sent through an object. The objects labeled 1 and 2 are described by the impulse response operators $\hat{h}_1$ and $\hat{h}_2$, respectively. The claim by Abouraddy *et al.* is that, if the source emits an entangled two-photon multimode state, and if one of the photons emitted is detected by a "bucket detector" (to be defined below), information about a test object is obtained that cannot be obtained with any classical source of light. Abouraddy *et al.* have proposed two schemes, in the first, called distributed quantum imaging, the test object, that we want to characterize, is described by $\hat{h}_1$, while $\hat{h}_2$ is a (known) reference object [9]. The authors have also suggested a scheme where the roles of the objects 1 and 2 are reversed [10], so that the test object is described by $\hat{h}_2$ and the photon that goes through it is detected by a bucket detector, while the photon going through the reference object, now described by $\hat{h}_1$, is detected by a detector array. This imaging method was named quantum holography.

The claim by Abouraddy *et al.* has been put in doubt by a recent analysis, and an experiment, by Bennink *et al.* [11]. Their analysis showed that if the test object in a quantum holographic setup is lossless, no information about the test object is obtained. In their experiment they used a lossy test object and a correlated classical source of light. Yet, using what Bennink *et al.* called "'two-photon' coincidence imaging" (where the authors' quotation marks indicate that the detection method mimics what Abouraddy *et al.* call quantum holography, but that classically correlated states were used instead of two-photon states) the information stored in the test object could be read out. In this paper, we extend the results of Bennink *et al.* by showing that distributed quantum imaging with a lossless reference object gives identically the same information about the test object as if an appropriate classical source were used. We also show that quantum holography with a lossless reference object (as in the experiment by Bennink *et al.*) can always be mimicked by a classical source.

We shall assume that the considered modes, both on the source side and on the detection side, are discrete. Apart from this difference, we shall try to keep the notation as similar as possible to the notation in Ref. [9]. Moreover, we will, just as Abouraddy *et al.* [9], assume that the two photons are spatially filtered in such a way that the photon emitted into unprimed modes does not interact with object 2, and the photon emitted into primed modes does not interact with object 1.

Consider a source that emits light in the pure two-photon state

$$|\Psi\rangle = \sum_i \sum_{j'} \varphi(i, j')|1_i, 1_{j'}\rangle, \tag{1}$$

where we have used the shorthand notation $|1_i\rangle \otimes |1_{j'}\rangle \equiv |1_i, 1_{j'}\rangle$. Let us first see what happens if we completely ignore the photons passing through object 2. Writing $\hat{\rho} = |\Psi\rangle\langle\Psi|$, the state of the unprimed photon then becomes

$$\mathrm{Tr}_2(\hat{\rho}) = \sum_{q'} \langle 1_{q'}|\hat{\rho}|1_{q'}\rangle. \tag{2}$$

In general, this is a mixed state of a single photon. After



FIG. 1: The two-photon source emits one photon into the modes denoted i, which propagate through object 1 described by the transformation $\hat{h}_1$. The linear impulse response operator $\hat{h}_1$ contains the information about object 1. Another photon is emitted into modes j' and is subjected to the operator $\hat{h}_2$ describing object 2. On the output side, there is an array of detectors, each measuring the photon number of a specific output mode.

interaction with object 1, the output state becomes

$$\hat{h}_1 \sum_{q'} \langle 1_{q'} | \hat{\rho} | 1_{q'} \rangle \hat{h}_1^\dagger = \sum_{q'} \langle 1_{q'} | \hat{h}_1 \hat{\rho} \hat{h}_1^\dagger | 1_{q'} \rangle, \tag{3}$$

where the equality follows from the fact that we have assumed that $\hat{h}_1$ does not operate on the primed vectors. It may be prudent, at this point, to establish that the mode sets on the input and output side of object 1 or 2 do not have to be of the same type, as long as both are enumerable and orthonormal. For any given object, every specific pair of mode sets will lead to a unique transformation $\hat{h}_1$. On the detector side it is convenient to work in the eigenmodes of the detectors. Then, the probability $p(q)$ of a photon detection in mode (detector) q is

$$p_1(q) = \sum_{q'} \langle 1_q, 1_{q'} | \hat{h}_1 \hat{\rho} \hat{h}_1^\dagger | 1_q, 1_{q'} \rangle. \tag{4}$$

If one defines $\gamma_1(i,j) = \sum_{k'} \varphi(i,k')\varphi^*(j,k')$, the coefficients $\gamma_1(i,j)$ turn out to be the density operator coefficients $\langle 1_i | \mathrm{Tr}_2(\hat{\rho}) | 1_j \rangle$, so that

$$\mathrm{Tr}_2(\hat{\rho}) = \sum_i \sum_j \gamma_1(i,j) |1_i\rangle\langle 1_j|. \tag{5}$$

By inserting the resolution of the identity twice into Eq. (4), one arrives at

$$p_1(q) = \sum_i \sum_j \gamma_1(i,j) h_1(q,i) h_1^*(q,j), \tag{6}$$



FIG. 2: Schematic drawing of "bucket detection" of the photon emitted into the primed modes corresponding to the lower detector array in the figure.

where $h_1(q,i) \equiv \langle 1_q | \hat{h}_1 | 1_i \rangle$ and $h_1^*(q,j) \equiv (\langle 1_q | \hat{h}_1 | 1_j \rangle)^\dagger$. The expression above is the direct discrete-mode equivalent to Eq. (8) in Ref. [9]. If one permutes the indices q and q', and lets the index $1 \to 2$ in Eqs. (4) and (6), one obtains the expressions for $p_2(q')$.

Now, assume that both photons in the photon pair are considered. The output state after the interaction with the two objects becomes $\hat{h}_1 \hat{h}_2 \hat{\rho} \hat{h}_2^\dagger \hat{h}_1^\dagger$. Note that, since $\hat{h}_1$ and $\hat{h}_2$ operate in different vector spaces, they commute. The probability of registering a correlated detection event between detector q and detector q' can be expressed as

$$p(q,q') = \langle 1_q, 1_{q'} | \hat{h}_1 \hat{h}_2 \hat{\rho} \hat{h}_2^\dagger \hat{h}_1^\dagger | 1_q, 1_{q'} \rangle. \tag{7}$$

A "bucket detector" is a multimode detector where all the modes propagating through an object are measured jointly. Hence, the information about the location of the detected photon (or equivalently, in what mode the photon was detected) is "erased." A schematic of such a detector is depicted in Fig. 2. In this case, the probability of photon detection in mode q (the marginal probability distribution of the photons interacting with object 1) becomes

$$\bar{p}_1(q) = \sum_{q'} \langle 1_q, 1_{q'} | \hat{h}_1 \hat{h}_2 \hat{\rho} \hat{h}_2^\dagger \hat{h}_1^\dagger | 1_q, 1_{q'} \rangle. \tag{8}$$

If we assume that the state from the source is entangled such that $\varphi(i,j) = \varphi(i)\delta_{ij'}$, Eq. (8) can be rewritten as

$$\bar{p}_1(q) = \sum_i \sum_j \sum_{q'} \varphi(i)\varphi^*(j) h_1(q,i) h_1^*(q,j) h_2(q',i) h_2^*(q',j). \tag{9}$$

Following Ref. [9], we also introduce the parameter $g_2(k',l') = \sum_{q'} h_2(q',k') h_2^*(q',l')$. Replacing the sum

over q' in Eq. (9) with this expression, we can write

$$\bar{p}_1(q) = \sum_i \sum_j \varphi(i)\varphi^*(j) g_2(i,j) h_1(q,i) h_1^*(q,j). \quad (10)$$

This is the discrete counterpart of Eq. (12c) in Ref. [9]. Comparing expressions (6) and (8), Abouraddy *et al.* claim that "based on classical probability analysis one would intuitively expect that $p_1(q)$ would be equal to $\bar{p}_1(q)$. This is not always the case, however." From a strictly mathematical point of view it is evident that $p_1(q)$ and $\bar{p}_1(q)$ are not equal for general functions $h_1(q,i)$ and $h_2(q',i')$. However, we shall now prove that if the reference object is lossless, the two expressions are equal.

Assume, therefore, that object 2, described by $\hat{h}_2$ is lossless. This immediately implies that *the operator* $\hat{h}_2$ *is unitary.* Using this fact, we can write the expression for $p_1(q)$ as

$$p_1(q) = \sum_{q'} \langle 1_q, 1_{q'} | \hat{h}_1 \hat{h}_2^\dagger \hat{h}_2 \hat{\rho} \hat{h}_2^\dagger \hat{h}_2 \hat{h}_1^\dagger | 1_q, 1_{q'} \rangle. \quad (11)$$

Furthermore, since $\hat{h}_1$ and $\hat{h}_2$ commute, we can recast this equation as

$$p_1(q) = \sum_{q'} \langle 1_q, 1_{q'} | \hat{h}_2^\dagger \hat{h}_1 \hat{h}_2 \hat{\rho} \hat{h}_2^\dagger \hat{h}_1^\dagger \hat{h}_2 | 1_q, 1_{q'} \rangle. \quad (12)$$

Since the trace operation is basis invariant, the trace Tr_2 of any operator $\hat{O}$ can be expressed either $\sum_{q'} \sum_{n=0}^{\infty} \langle n_{q'} | \hat{O} | n_{q'} \rangle$ or $\sum_{q'} \sum_{n=0}^{\infty} \langle n_{q'} | \hat{h}_2^\dagger \hat{O} \hat{h}_2 | n_{q'} \rangle$, where $\hat{h}_2$ is an arbitrary, unitary operator. Hence, from Eq. (12), we get

$$p_1(q) = \sum_{q'} \langle 1_q, 1_{q'} | \hat{h}_1 \hat{h}_2 \hat{\rho} \hat{h}_2^\dagger \hat{h}_1^\dagger | 1_q, 1_{q'} \rangle = \bar{p}_1(q). \quad (13)$$

This relation implies that Eqs. (4) and (8) are *identical* if the object 2 (the reference object) is lossless, irrespective of $\hat{\rho}$ and $\hat{h}_1$. Hence, in this case, nothing is gained by using distributed quantum imaging over single photon (per necessity, uncorrelated) imaging. The single-photon state $\mathrm{Tr}_2(\hat{\rho})$ will give the same detection statistics as $\hat{\rho}$.

Let us now consider the case where the test object is described by $\hat{h}_2$, that is, it is the modes through the test object that are detected with the "bucket detector." This imaging principle is called "quantum holography" [10] or "two-photon coincidence imaging" [11]. Abouraddy *et al.* [10] claimed that it is possible to use entanglement between the states in the primed and unprimed modes to read out holographic information about the test object $\hat{h}_2$ even if the photons traversing it are detected with a bucket detector, such as a sphere coated on the inside with a photosensitive film that surrounds it. However, Bennink *et al.* showed that if the test object is lossless, then the bucket detector always clicks, effectively tracing out all information about the test object. Hence, quantum holography of such test objects does not work.

However, assume that the test object (object 2) is lossy. It is clear that in this case the reference object (object 1) should be lossless, as losses will only introduce additional randomness in the measurement. Since $\hat{h}_1$ and $\hat{h}_2$ commute, this randomness corresponds to loss of information about $\hat{h}_2$. If the reference object is lossless, it is rather obvious that if a two-photon (possibly) entangled state $|\Psi\rangle$ is used, its quantum holography detection statistics will be exactly mimicked by the classically correlated state

$$\hat{\rho}' = \sum_i \hat{h}_1^\dagger |1_i\rangle\langle 1_i| \hat{h}_1 \otimes \langle 1_i | \hat{h}_1 \hat{\rho} \hat{h}_1^\dagger | 1_i \rangle. \quad (14)$$

Hence, it seems to us that entangled-state holography with bucket detection *of any object* is of limited use, since identical results can be obtained using classical states. This was what Bennink *et al.* [11] demonstrated experimentally. The statistical distribution $\bar{P}_1(q)$ in their experiment very clearly brought out the information encoded in the mask although all the photons interacting with the (lossy) mask were detected by a bucket detector.

Above, we have shown that distributed quantum imaging and quantum holography with a lossless reference object (but, in general, a lossy test object) will not offer anything imaging with classical states cannot provide. Bennink *et al.* [11] proved that quantum holography of a lossless test object also can be mimicked by classical states. Now, we will treat the case when both object 1 and 2 are lossy. The standard way of including losses is to extend the two original sets of modes with auxiliary modes. We shall do so by assuming that the respective detector eigenmodes constitute only a subset of the primed and unprimed modes. To be realistic we can assume that only the photons in N (N') of the unprimed (primed) modes are detected, and that these modes are labeled $1, 2, \ldots, N$ ($1', 2', \ldots, N'$). The probability of detecting a photon in mode q then becomes

$$\begin{aligned} P_1(q) &= \sum_{q'=1}^{N'} \langle 1_q, 1_{q'} | \hat{h}_1 \hat{h}_2 \hat{\rho} \hat{h}_2^\dagger \hat{h}_1^\dagger | 1_q 1_{q'} \rangle \\ &\quad + \sum_{q'>N'} \langle 1_q, 1_{q'} | \hat{h}_1 \hat{h}_2 \hat{\rho} \hat{h}_2^\dagger \hat{h}_1^\dagger | 1_q, 1_{q'} \rangle \\ &= \bar{P}_1(q) + P_{1,0}(q). \end{aligned} \quad (15)$$

From Eq. (15), we see that photodetection probability $P_1(q)$ is simply the sum over all the detection events q' of the two-photon joint probability distribution $\bar{P}_1(q) = \sum_{q'=1}^{N'} P(q,q')$ and the probability of detecting one photon in mode q and no photon in the modes $q' = 1', 2', \ldots, N'$. This is just what one expects intuitively. For lossy reference objects, $P_1(q)$ differs from $\bar{P}_1(q)$.

Now, denote the probability of not detecting the photon in the primed modes $P_0 = \sum_q P_{1,0}(q)$ and consider,

e.g., the product state

$$\sum_{j'=1'}^{N'} \langle 1_{j'}|\hat{h}_2\hat{\rho}\hat{h}_2^\dagger|1_{j'}\rangle \otimes \hat{h}_2^\dagger \left(|1_{1'}\rangle\langle 1_{1'}| + \frac{P_0|1_{M'}\rangle\langle 1_{M'}|}{1-P_0} \right) \hat{h}_2, \tag{16}$$

where $M' > N'$. It is quite obvious that this state, that lacks any correlation between the photon in the primed and unprimed modes, will yield an identical photodetection statistics distribution $\bar{P}_1(q)$ to the corresponding distribution of the state $\hat{\rho}$ (that may be entangled). Note that this is true irrespective of $\hat{\rho}$, $\hat{h}_1$, and $\hat{h}_2$. The result depends critically on the fact that $\hat{h}_2$ is an operation local in the primed modes. However, this is, in general, not a physically accessible state since it may contain excitation in modes with $M' > N'$ (the modes corresponding to the loss). If only the modes $q' = 1', \ldots, N'$ can be excited, then one cannot, in general, mimic the photodetection statistics of the two lossy objects illuminated by an two-photon entangled state, with a classically correlated state. However, even if this is not the case, we do not see what could be gained by using entanglement, because, in general, loss will destroy the entanglement of a quantum state more rapidly than it will reduce the correlations in a classical state. Since we have shown that in the case when the reference object is lossless, entanglement combined with bucket detection offers no improvement for neither two-photon imaging nor for quantum holography, we conjecture that *if bucket detection of one of the photons is used*, entangled two-photon imaging never offers any advantage over (classically) correlated-photon imaging.

An effect similar to "quantum holography," but not to be confused with it, has been demonstrated in Refs. [12, 13], where a diffraction pattern could be read out trough the joint probability distribution of a two-photon state, although the marginal probability distributions did not show any diffraction pattern. This is an effect due to entanglement and the fact that *bucket detection was not used.* To model such an effect in the simplest possible fashion, consider the two-photon, four-mode, entangled state

$$|\Psi\rangle = \frac{1}{2}(|1_1, 1_{1'}\rangle + |1_1, 1_{2'}\rangle + |1_2, 1_{1'}\rangle - |1_2, 1_{2'}\rangle). \tag{17}$$

Assume that $\hat{h}_1 = \hat{1}_1$ and $h_2(1,1) = h_2(1,2) = h_2(2,1) = -h_2(2,2) = 1/\sqrt{2}$. One can then readily show that $p(1,1') = p(2,2') = 1/2$ and $p(1,2') = p(2,1') = 0$. That is, which one of the (two) detectors behind each object that will "click" is totally unpredictable. The statistics of either detector pair alone do not contain any information about object 2. This is so regardless if one simply ignores the outcome of the other detector pair, or if one uses a bucket detector (that, in this case, always will "click"). Due to the input-state entanglement, there is, however, a perfect correlation between the joint detection events. The correlation statistics hence reveal (some) information about the object.

In summary, we have shown that although entangled two-photon imaging may bring out effects that cannot be mimicked by any classical source, the advantage with this imaging method is lost if "bucket detection" of one of the photons is employed, at least if the reference object is lossless (and this ideal case is what one should strive for, since it gives the optimal information about the test object). To capitalize on the unique properties of two-photon imaging, the detection joint probability distribution needs to be retained.

The authors would like to thank Professor B. E. A. Saleh for useful comments on this work. The work was supported by the Swedish Research Council (VR) and the Swedish Foundation for Strategic Research (SSF).

---

* Electronic address: gunnarb@imit.kth.se; URL: `http://www.ele.kth.se/QEO/`

[1] M. I. Kolobov and I. V. Sokolov, Phys. Lett. A **140**, 101 (1989).

[2] M. I. Kolobov and C. Fabre, Phys. Rev. Lett. **85**, 3789 (2000).

[3] C. Fabre, J. B. Fouet, and A. Maître, Opt. Lett. **25**, 76 (2000).

[4] A. N. Boto, P. Kok, D. S. Abrams, S. L. Braunstein, C. P. Williams, and J. P. Dowling, Phys. Rev. Lett. **85**, 2733 (2000).

[5] P. Kok, A. N. Boto, D. S. Abrams, C. P. Williams, S. L. Braunstein, and J. P. Dowling, Phys. Rev. A **63**, 063407 (2001).

[6] G. Björk, L. L. Sánchez-Soto, and J. Söderholm, Phys. Rev. Lett. **86**, 4516 (2001).

[7] G. Björk, L. L. Sánchez-Soto, and J. Söderholm, Phys. Rev. A **64**, 013811 (2001).

[8] J. Jacobson, G. Björk, I. Chuang, and Y. Yamamoto, Phys. Rev. Lett. **74**, 4835 (1995).

[9] A. F. Abouraddy, B. E. A. Saleh, A. V. Sergienko, and M. C. Teich, Phys. Rev. Lett. **87**, 123602 (2001).

[10] A. F. Abouraddy, B. E. A. Saleh, A. V. Sergienko, and M. C. Teich, Opt. Express **9**, 498 (2001).

[11] R. S. Bennink, S. J. Bentley, and R. W. Boyd, Phys. Rev. Lett. **89**, 113601 (2002).

[12] D. V. Strekalov, A. V. Sergienko, D. N. Klyshko, and Y. H. Shih, Phys. Rev. Lett. **74**, 3600 (1995).

[13] T. B. Pittman, Y. H. Shih, D. V. Strekalov, and A. V. Sergienko, Phys. Rev. A **52**, R3429 (1995).